arXiv:1608.06377v2 [hep-th] 19 Sep 2016

Hawking Radiation of Vector Particles via Tunneling From 4-Dimensional And 5-Dimensional Black Holes

Zhongwen Feng, [1] * Yang Chen, [2] Xiaotao Zu [1] †

1. School of Physical Electronics, University of Electronic Science and Technology of China, Chengdu, 610054, China

2. Institute of Theoretical Physics, China West Normal University, Nanchong, 637009, China

Abstract: Using Proca equation and WKB approximation, we investigate Hawking radiation of vector particles via tunneling from 4-dimensional Kerr-de Sitter black hole and 5-dimensional Schwarzschild-Tangherlini black hole. The results show that the tunneling rates and Hawking temperatures are depended on the properties of spacetime (event horizon, mass and angular momentum). Besides, our results are the same as scalars and fermions tunneling from 4-dimensional Kerr-de Sitter black hole and 5-dimensional Schwarzschild-Tangherlini black hole.

Keywords: Vector particles; Hawking temperature; Quantum tunneling

1 Introduction

In 1970s, Bekenstein and Hawking discovered the thermodynamic of black holes, which indicates that black holes have thermal radiation (Bekenstein, 1973; Hawking, 1975). Later, Hawking investigated the radiation from black holes by quantum mechanics, and presented the theory of Hawking radiation. The existence of thermodynamic for black hole was an important discovery, which great influence on the foundations of physics. Thus, the Hawking radiation attracted researchers' attention.

The quantum tunneling for black holes is considered as an important method to study the Hawking radiation. This method was put forward by Parikh and Wilczek (2000). Later, Kerner and Mann (2008a); Kerner and Mann (2008b) developed the quantum tunneling method and studied the Dirac particles tunneling from spherically symmetric black holes. The Hamilton-Jacobi ansatz is another kind of tunneling method. Using the WKB approximation, the tunneling rate can be calculated by the formula $\Gamma \propto \exp\left(-2\mathrm{Im}S_0\right)$, where S_0 is the classical action at the leading order in $\hbar$, then the Hawking temperature is obtained (Angheben et al., 2005; Shankaranarayanan, 2005). Subsequently, higher order calculations of scalars, fermions and bosons tunneling from black holes are studied by Wang et al. (2010); Chatterjee and Mitra (2009); Yale (2011a,b). By defining a new of the particle energy, their work is consistent with the original results, which incidents there is no higher-order corrections to the Hawking temperature. Following that, a lot of work has been done for studying the Hawking radiation from black holes (Feng et al., 2012; Jiang et al., 2006; Lin and Yang , 2014; Chen et al., 2013a,b; Chen and Li, 2014; Zhang and Zhao , 2005c; Jiang, 2008; Ding and Jing, 2009; Zhang and Zhao, 2005a,b, 2012; Li et al., 2012; Li and Lin, 2011; Mu et al., 2010).

Obviously, the black holes do not only radiate scalar particles and Dirac particles, they may emit particles with arbitrary spin. Recently, Kruglov (2014b,a) and Chen et al. (2015) researched the vector (namely, spin-1 bosons) tunneling from black holes. However, all of their work are limited to the vector particles tunneling from the low dimensional (1+1 dimensional black hole and 1+2 dimensional) black holes. In this paper, the vector particles tunneling from 4-dimensional Kerr-de Sitter (KdS) black hole and 5-dimensional Schwarzschild-Tangherlini (ST) black hole are investigated with the help of the Proca equation and WKB approximation. The KdS black hole is an important rotating solution of the Einstein equation with a positive cosmological constant. As we know, the cosmological constant is one kind of candidates for dark energy, it consistent with the current Lambda-CDM original model, and can explain the accelerating expansion of our universe. Besides, the KdS black holes have four horizons. Its special structure can help people further understand the properties of gravity. On the other hand, the 5-dimensional ST black hole is a good approximation to a 5 dimensional compactified black hole, people can use it to study the higher dimensional distorted compactified spacetime. From what has been discussed above, we think the vector tunneling from the 4-dimensional KdS black hole and the 5-dimensional ST black hole are worth to be studied, people may obtain more information of Hawking radiation from the higher-dimensional black holes.

*E-mail: zwfengphy@163.com

†E-mail: xtzu@uestc.edu.cn

The paper is organized as follows. In next section, extending the work of Kruglov (2014a), the vector tunneling from 4-dimensional KdS black hole will be investigated. In section 3, in the 5-dimensional ST spacetime, the spin-1 bosons tunneling radiation is derived. Section 4 is devoted to our discussions and conclusions.

2 Vector tunneling from the 4-dimensional Kerr-de Sitter black hole

Extending the method which presented by Kruglov, the vector tunneling from the 4-dimensional KdS black hole is investigated in this section. In Carter (1968), the metric of 4-dimensional KdS black hole is given by

$$ds^2 = -\frac{\Delta}{\rho^2}\left(dt - \frac{a\sin^2\theta}{\Xi}d\phi\right)^2 + \rho^2\left(\frac{1}{\Delta}dr^2 + \frac{1}{\Delta_\theta}d\theta^2\right) + \frac{\Delta_\theta\sin^2\theta}{\rho^2}\left(adt - \frac{r^2+a^2}{\Xi}d\phi\right)^2, \tag{1}$$

where $\Delta = \left(r^2+a^2\right)\left(1-r^2l^{-2}\right) - 2Mr$, $\Delta_\theta = 1 + a^2l^{-2}\cos^2\theta$, $\Xi = 1 + a^2l^{-2}$ and $\rho^2 = r^2 + a^2\cos^2\theta$, M and a are the mass and angular momentum of 4-dimensional KdS black hole, l is a constant depended on the cosmological factor as $\Lambda = 3l^{-2}$, respectively. One can obtains four horizons when $\Delta = 0$. The four horizons are outer cosmological horizon r_{c+}, outer event horizon r_+, inner event horizon r_- and inner cosmological horizon r_{c-}, which satisfy the relation $r_{c+} > r_+ > r_- > 0 > r_{c-}$. For convenience, we redefined Eq.(1) as

$$ds^2 = -A(r)dt^2 + B(r)^{-1}dr^2 + C(r)d\theta^2 + D(r)d\phi^2 + 2E(r)dtd\phi, \tag{2}$$

the notations are $A(r) = \Delta - a\Delta_\theta\sin^2\theta/\rho^2$, $B(r) = \Delta/\rho^2$, $C(r) = \rho^2/\Delta_\theta$, $D(r) = \sin^2\theta[\Delta_\theta(a^2+r^2)^2 - a^2\Delta\sin^2\theta]/\Xi^2\rho^2$, and $E(r) = a\sin^2\theta[\Delta + (a^2+r^2)]\Delta_\theta/\Xi\rho$, respectively.

According to Kruglov (2014b,a, 2011), the dynamic behavior of vector particles in curved spacetime is described by Proca equation

$$D_\mu\psi^{\nu\mu} + \frac{m^2}{\hbar^2}\psi^\nu = 0, \tag{3}$$

$$\psi_{\nu\mu} = D_\nu\psi_\mu - D_\mu\psi_\nu = \partial_\nu\psi_\mu - \partial_\mu\psi_\nu, \tag{4}$$

where D_μ are covariant derivatives, ψ_ν is related to the ψ_t, ψ_r, ψ_θ and ψ_ϕ, m is the mass of vector particles. $\psi^{\mu\nu}$ is an anti-symmetrical tensor. Thus, the Eq. (3) can be rewritten as follows

$$\frac{1}{\sqrt{-g}}\partial_\mu\left(\sqrt{-g}\psi^{\nu\mu}\right) + \frac{m^2}{\hbar^2}\psi^\nu = 0. \tag{5}$$

The components of ψ^ν and $\psi^{\mu\nu}$ are

$$\begin{aligned}
&\psi^0 = \tfrac{-D\psi_0+E\psi_3}{AD+E^2}, \quad \psi^1 = B\psi_1, \quad \psi^2 = C^{-1}\psi_2,\\
&\psi^3 = \tfrac{E\psi_0+A\psi_3}{AD+E^2}, \quad \psi^{01} = \tfrac{-DB\psi_{01}-EB\psi_{13}}{AD+E^2},\\
&\psi^{02} = \tfrac{C^{-1}(-D\psi_{02}-E\psi_{23})}{AD+E^2}, \quad \psi^{03} = -\tfrac{\psi_{03}}{AD+E^2},\\
&\psi^{12} = BC^{-1}\psi_{12}, \quad \psi^{13} = \tfrac{B(A\psi_{13}-E\psi_{01})}{AD+E^2},\\
&\psi^{23} = \tfrac{C^{-1}(A\psi_{23}-E\psi_{02})}{AD+E^2}.
\end{aligned} \tag{6}$$

Substituting Eq. (4) and Eq. (6) into Eq. (3), one yields

$$\begin{aligned}
&\tfrac{1}{\sqrt{-g}}\left\{\partial_r\left[\sqrt{-g}\left(\tfrac{-DB\psi_{01}-EB\psi_{13}}{AD+E^2}\right)\right] + \partial_\theta\left[\sqrt{-g}\left(\tfrac{-DC^{-1}\psi_{02}-EC^{-1}\psi_{23}}{AD+E^2}\right)\right] +\right.\\
&\left.\partial_\phi\left[\sqrt{-g}\left(-\tfrac{\psi_{03}}{AD+E^2}\right)\right]\right\} + \tfrac{m^2}{\hbar^2}\left(\tfrac{E\psi_3-D\psi_0}{AD+E^2}\right) = 0,
\end{aligned} \tag{7}$$

$$\begin{aligned}&\tfrac{1}{\sqrt{-g}}\left\{\partial_t\left[\sqrt{-g}\left(\tfrac{DB\psi_{01}+EB\psi_{13}}{AD+E^2}\right)\right]+\partial_\theta\left(\sqrt{-g}BC^{-1}\psi_{12}\right)\right.\\&\left.+\partial_\phi\left[\sqrt{-g}\left(\tfrac{AB\psi_{13}-BE\psi_{01}}{AD+E^2}\right)\right]\right\}+\tfrac{m^2}{\hbar^2}B\psi_1=0,\end{aligned}\tag{8}$$

$$\begin{aligned}&\tfrac{1}{\sqrt{-g}}\left\{\partial_t\left[\sqrt{-g}\left(\tfrac{DC^{-1}\psi_{02}+EC^{-1}\psi_{23}}{AD+E^2}\right)\right]+\partial_\theta\left[\sqrt{-g}\left(-BC^{-1}\psi_{12}\right)\right]\right.\\&\left.+\partial_\phi\left[\sqrt{-g}\left(\tfrac{C^{-1}A\psi_{23}-C^{-1}E\psi_{02}}{AD+E^2}\right)\right]\right\}+\tfrac{m^2}{\hbar^2}C^{-1}\psi_2=0,\end{aligned}\tag{9}$$

$$\begin{aligned}&\tfrac{1}{\sqrt{-g}}\left\{\partial_t\left[\sqrt{-g}\left(\tfrac{\psi_{03}}{AD+E^2}\right)\right]+\partial_r\left[\sqrt{-g}\left(\tfrac{BE\psi_{01}-AB\psi_{13}}{AD+E^2}\right)\right]\right.\\&\left.+\partial_\theta\left[\sqrt{-g}\left(\tfrac{C^{-1}E\psi_{02}-C^{-1}A\psi_{23}}{AD+E^2}\right)\right]\right\}+\tfrac{m^2}{\hbar^2}\left(\tfrac{E\psi_0+A\psi_3}{AD+E^2}\right)=0.\end{aligned}\tag{10}$$

For solving Eqs. (7) - (10), ψ_ν taking form as

$$\psi_\nu=(c_\nu)\exp\left[\frac{i}{\hbar}S_0\left(t,r,\theta,\phi\right)+\sum_n\hbar^nS_n\left(t,r,\theta,\phi\right)\right],\tag{11}$$

where $n=1,2,3,\cdots$. Since the WKB approximation is been applied here, the higher order terms of $\mathcal{O}(\hbar)$ are neglected. The resulting equations to leading order in $\hbar$ are

$$\begin{aligned}&DB\left[c_1\left(\partial_tS\right)\left(\partial_rS\right)-c_0\left(\partial_rS\right)^2\right]+EB\left[c_3\left(\partial_rS\right)^2-c_1\left(\partial_\phi S\right)\left(\partial_rS\right)\right]\\&+DC^{-1}\left[c_2\left(\partial_tS\right)\left(\partial_\theta S\right)-c_0\left(\partial_\theta S\right)^2\right]+EC^{-1}\left[c_3\left(\partial_\theta S\right)^2-c_2\left(\partial_\phi S\right)\left(\partial_\theta S\right)\right]\\&+\left[c_3\left(\partial_\phi S\right)\left(\partial_tS\right)-c_0\left(\partial_\phi S\right)^2\right]+m^2\left(c_3E-c_0D\right)=0,\end{aligned}\tag{12}$$

$$\begin{aligned}&DB\left[c_0\left(\partial_rS\right)\left(\partial_tS\right)-c_1\left(\partial_tS\right)^2\right]+EB\left[c_3\left(\partial_tS\right)\left(\partial_rS\right)c_1\left(\partial_\phi S\right)\left(\partial_tS\right)\right]\\&+BC^{-1}\left(AD+E^2\right)\left[c_1\left(\partial_\theta S\right)^2-c_2\left(\partial_\theta S\right)\left(\partial_rS\right)\right]+BE\left[c_1\left(\partial_tS\right)\left(\partial_\varphi S\right)-c_0\left(\partial_rS\right)\left(\partial_\phi S\right)\right]\\&+AB\left[c_1\left(\partial_\phi S\right)^2-c_3\left(\partial_rS\right)\left(\partial_\phi S\right)\right]+c_1m^2B\left(AD+E^2\right)=0,\end{aligned}\tag{13}$$

$$\begin{aligned}&DC^{-1}\left[c_0\left(\partial_tS\right)\left(\partial_\theta S\right)-c_2\left(\partial_tS\right)^2\right]+EC^{-1}\left[c_3\left(\partial_tS\right)\left(\partial_\theta S\right)-c_2\left(\partial_tS\right)\left(\partial_\phi S\right)\right]\\&+BC^{-1}\left(AD+E^2\right)\left[c_2\left(\partial_rS\right)^2-c_1\left(\partial_rS\right)\left(\partial_\theta S\right)\right]+C^{-1}E\left[c_2\left(\partial_tS\right)\left(\partial_\phi S\right)-c_0\left(\partial_\phi S\right)\left(\partial_\theta S\right)\right]\\&+C^{-1}A\left[c_2\left(\partial_\phi S\right)^2-c_3\left(\partial_\theta S\right)\left(\partial_\phi S\right)\right]+c_2m^2C^{-1}\left(AD+E^2\right)=0,\end{aligned}\tag{14}$$

$$\begin{aligned}&\left[c_0\left(\partial_\phi S\right)\left(\partial_tS\right)-c_3\left(\partial_tS\right)^2\right]+BE\left[c_0\left(\partial_rS\right)^2-c_1\left(\partial_tS\right)\left(\partial_rS\right)\right]\\&+AB\left[c_1\left(\partial_rS\right)\left(\partial_\phi S\right)-c_3\left(\partial_rS\right)^2\right]+C^{-1}E\left[c_0\left(\partial_\theta S\right)^2-c_2\left(\partial_\theta S\right)\left(\partial_tS\right)\right]\\&+C^{-1}A\left[c_2\left(\partial_\theta S\right)\left(\partial_\phi S\right)-c_3\left(\partial_\theta S\right)^2\right]+m^2\left(c_0E+c_3A\right)=0.\end{aligned}\tag{15}$$

Considering the spacetime of metric (1) has two Killing vectors ∂_t and ∂_ϕ, the solutions of Eqs. (12) - (15) are in the form

$$S_0 = -\omega t + W(r) + \Theta(\theta) + j\phi, \tag{16}$$

where ω and j are the energy and the angular momentum of vector particles, respectively. Putting Eq. (16) into Eqs. (12) - (15), one gets

$$\Lambda (c_0, c_1, c_2, c_3)^T = 0. \tag{17}$$

The Λ in Eq. (17) is a 4×4 matrix, its components are

$$\begin{aligned}
&\Lambda_{00} = -Dm^2 - j^2 - BD(W')^2 - DC^{-1}(\partial_\theta\Theta)^2, \Lambda_{01} = -BEW'j - BDW'\omega,\\
&\Lambda_{02} = -C^{-1}Ej(\partial_\theta\Theta) - C^{-1}D(\partial_\theta\Theta)\omega, \Lambda_{03} = BE(W')^2 + Em^2 + C^{-1}E(\partial_\theta\Theta)^2 - j\omega,\\
&\Lambda_{10} = -BEW'(\partial_\theta\Theta) - BDW'\omega, \Lambda_{11} = m^2B\left(AD + E^2\right) - DB\omega^2 - EBj\omega - C^{-1}B(\partial_\theta\Theta)^2 - EB(\partial_\theta\Theta)\omega - ABj^2,\\
&\Lambda_{12} = -BC^{-1}W'(\partial_\theta\Theta)\left(AD + E^2\right), \Lambda_{13} = BEW'\omega - ABW'j, \Lambda_{20} = EC^{-1}j(\partial_\theta\Theta) - C^{-1}D(\partial_\theta\Theta)\omega,\\
&\Lambda_{21} = -BC^{-1}(\partial_\theta\Theta)W'\left(AD + E^2\right), \Lambda_{22} = m^2C^{-1}\left(AD + E^2\right) - C^{-1}D\omega^2 - C^{-1}Ej\omega\\
&-BC^{-1}(W')^2 - AC^{-1}j^2 - C^{-1}EjW', \Lambda_{23} = C^{-1}E(\partial_\theta\Theta)\omega - A(r)C(r)^{-1}j(\partial_\theta\Theta),\\
&\Lambda_{30} = m^2E - j\omega + BE(W')^2 + EC^{-1}(\partial_\theta\Theta)^2, \Lambda_{31} = BEW'\omega - ABjW',\\
&\Lambda_{32} = EC^{-1}(\partial_\theta\Theta)\omega - AC^{-1}j\Theta, \Lambda_{33} = Am^2 - \omega^2 + AB(W')^2 + AC^{-1}(\partial_\theta\Theta)^2,
\end{aligned} \tag{18}$$

where $W' = \partial_r S_0$ and $j = \partial_\phi S_0$. Eq. (17) has a nontrivial solution when $\mathrm{Det}(\Lambda) = 0$. Hence, one yields

$$W_\pm = \int \sqrt{\frac{(\omega^2 - \Omega j)^2 + X}{(AD + E^2)BD^{-1}}} dr = \pm i\pi \frac{\omega - \Omega(r_+)j}{2\kappa(r_+)}, \tag{19}$$

where $+(-)$ are the outgoing (incoming) solutions on the outer event horizon, $X = -D^{-2}[E^2j^2 + m^2D(AD + E^2)] + C^{-1}D^{-1}[(AD + E^2) - AC]$, $\Omega(r_+) = -E(r_+)/D(r_+) = a\Xi/\left(r_+^2 + a^2\right)$ is the angular velocity on outer the event horizon and $\kappa(r_+) = r_+(1 - 2r_+^2l^{-2} - r_+a^2l^{-2})/(r_+^2 + a^2)$ is the surface gravity of outer event horizon. The tunneling rate of vector particle from 4-dimensional KdS black hole is obtained as

$$\Gamma = \frac{\Gamma_{(emission)}}{\Gamma_{(absorpation)}} = \frac{\exp(-2\mathrm{Im}W_+ - 2\mathrm{Im}\Xi)}{\exp(-2\mathrm{Im}W_- - 2\mathrm{Im}\Xi)}. \tag{20}$$

As we known, any particle outside the event horizon will fall into the black hole, one has $\Gamma_{absorpation} = 1$, namely, $\mathrm{ImW}_- + \mathrm{Im}\Xi = 0$. Therefore, the result is

$$\Gamma = \exp\left[-2\pi \frac{\omega - j\Omega(r_+)}{\kappa(r_+)}\right]. \tag{21}$$

With the help of Boltzman factor Feng et al. (2014), the Hawking temperature of 4-dimensional KdS black hole is

$$T_H = \frac{r_+ - r_+^3l^{-2} - r_+a^2l^{-2} - M}{2\pi\left(r_+^2 + a^2\right)}. \tag{22}$$

Eq. (21) and Eq. (22) are the vector particles tunneling rate and Hawking temperature of KdS black hole. The results showed that the Γ and T_H are depended on the outer event horizon r_+, mass M, angular momentum of KdS black hole a and the cosmological factor Λ. When $\Lambda = 0$, the Eq. (22) is reduced to the temperature of Kerr black hole. When $a = 0$, $\Lambda = 0$ and $r_+ = 2M$, the Hawking temperature of Schwarzschild (SC) black hole $T_{H(sc)} = 1/8\pi M$ is recovered (Zheng, 2008; Khani and Darvishi, 2008). Moreover, our result is fully in accordance with that obtained by other methods (Chen et al., 2013b; Li et al., 2012; Jiang, 2007; Wang et al., 2006; Ali et al., 2015).

3 Vector tunneling from the 5-dimensional Schwarzschild-Tangherlini black hole

In this section, we study the quantum tunneling from 5 dimensional Schwarzschild-Tangherlini black hole via Proca equation. When added extra compact spatial dimensions into a static spherically symmetric solution of the vacuum Einstein equations, one gets the line element of D-dimensional ST black hole Tangherlini (1963)

$$ds^2 = -f(r)\,dt^2 + f(r)^{-1}\,dr^2 + r^2 d\Omega_{D-2}^2, \tag{23}$$

where $f(r) = 1-(r_H/r)^{D-3}$, r_H marks the event horizon of the D-dimensional ST black hole, which is related to the mass of ST black hole as $M = (D-3)\,r_H^{D-3}/2$, $d\Omega_{D-2}^2 = d\chi_2^2 + \sin^2\chi_2 d\chi_3^2 + \cdots + \left(\prod_{k=2}^{D-2}\sin^2\chi_k\right) d\chi_{D-1}^2$ is the line element on $D-2$-sphere with the angles on this sphere χ_k. For $D=5$, one has Abdolrahimi et al. (2010)

$$\begin{aligned} ds^2 &= -\widetilde{A}(r)\,dt^2 + \widetilde{B}^{-1}(r)\,dr^2 + \widetilde{C}(r)\,d\zeta^2 + \widetilde{D}(r)\,d\vartheta^2 + \widetilde{E}(r)\,d\phi^2 \\ &= -\left(1-r_H^2/r^2\right)dt^2 + \left(1-r_H^2/r^2\right)^{-1}dr^2 + r^2 d\zeta^2 + r^2\sin^2\zeta d\vartheta^2 + r^2\sin^2\zeta\sin^2\vartheta d\phi^2, \end{aligned} \tag{24}$$

where $r_H^2 \sim M$ Beach (2014). Near the event horizon, it is clear that $\widetilde{B}(r) = \widetilde{B}'(r_H)(r-r_H) + \mathcal{O}\left[(r-r_H)^2\right]$. According to Eq. (4), one yields

$$\begin{aligned} &\psi^0 = -\widetilde{A}^{-1}\psi_0, \quad \psi^1 = \widetilde{B}\psi_1, \quad \psi^2 = \widetilde{C}^{-1}\psi_2, \\ &\psi^3 = \widetilde{D}^{-1}\psi_3, \quad \psi^4 = \widetilde{E}^{-1}\psi_4, \quad \psi^{01} = -\widetilde{B}\widetilde{A}^{-1}\psi_{01}, \\ &\psi^{02} = -(\widetilde{A}\widetilde{C})^{-1}\psi_{02}, \quad \psi^{03} = -(\widetilde{A}\widetilde{D})^{-1}\psi_{03}, \\ &\psi^{04} = -(\widetilde{A}\widetilde{E})^{-1}\psi_{04}, \quad \psi^{12} = \widetilde{B}\widetilde{C}^{-1}\psi_{12}, \\ &\psi^{13} = \widetilde{B}\widetilde{D}^{-1}\psi_{13}, \quad \psi^{14} = \widetilde{B}\widetilde{E}^{-1}\psi_{14}, \\ &\psi^{23} = (\widetilde{C}\widetilde{D})^{-1}\psi_{23}, \quad \psi^{24} = (\widetilde{C}\widetilde{E})^{-1}\psi_{24}, \\ &\psi^{34} = \left(\widetilde{D}\widetilde{E}\right)^{-1}\psi_{34}. \end{aligned} \tag{25}$$

Inserting Eq. (25) and $\psi_\nu = (c_\nu)\exp[\frac{i}{\hbar}S_0(t,r,\zeta,\vartheta,\phi)]$ into Eq. (3), and ignoring the higher order of terms of $\hbar$, we have

$$\begin{aligned} &\widetilde{B}\left[c_0(\partial_r S_0)^2 - c_1(\partial_r S_0)(\partial_t S_0)\right] + \widetilde{C}^{-1}\left[c_0(\partial_\zeta S_0)^2 - c_2(\partial_\zeta S_0)(\partial_t S_0)\right] + \\ &\widetilde{D}^{-1}\left[c_0(\partial_\vartheta S_0)^2 - c_3(\partial_\vartheta S_0)(\partial_t S_0)\right] + \widetilde{E}^{-1}\left[c_0(\partial_\phi S_0)^2 - c_4(\partial_\phi S_0)(\partial_t S_0)\right] \\ &+c_0 m^2 = 0, \end{aligned} \tag{26}$$

$$\begin{aligned} &\widetilde{A}^{-1}\left[c_0(\partial_r S_0)(\partial_t S_0) - c_1(\partial_t S_0)^2\right] + \widetilde{C}^{-1}\left[c_1(\partial_\zeta S_0)^2 - c_2(\partial_r S_0)(\partial_\zeta S_0)\right] \\ &+\widetilde{D}^{-1}\left[c_1(\partial_\vartheta S_0)^2 - c_3(\partial_r S_0)(\partial_\vartheta S_0)\right] + \widetilde{E}^{-1}\left[c_1(\partial_\phi S_0)^2 - c_4(\partial_r S_0)(\partial_\phi S_0)\right] \\ &+c_1 m^2 = 0, \end{aligned} \tag{27}$$

$$\widetilde{A}^{-1}\left[c_0\left(\partial_\zeta S_0\right)\left(\partial_t S_0\right)-c_2\left(\partial_t S_0\right)^2\right]+\widetilde{B}\left[c_2\left(\partial_r S_0\right)^2-c_1\left(\partial_\zeta S_0\right)\left(\partial_r S_0\right)\right]$$
$$+\widetilde{D}^{-1}\left[c_2\left(\partial_\vartheta S_0\right)^2-c_3\left(\partial_\zeta S_0\right)\left(\partial_\vartheta S_0\right)\right]+\widetilde{E}^{-1}\left[c_2\left(\partial_\phi S_0\right)^2-c_4\left(\partial_\zeta S_0\right)\left(\partial_\phi S_0\right)\right] \tag{28}$$
$$+c_2m^2=0,$$

$$\widetilde{A}^{-1}\left[c_0\left(\partial_\vartheta S_0\right)\left(\partial_t S_0\right)-c_3\left(\partial_t S_0\right)^2\right]+\widetilde{B}\left[c_3\left(\partial_r S_0\right)^2-c_1\left(\partial_r S_0\right)\left(\partial_\vartheta S_0\right)\right]$$
$$+\widetilde{C}^{-1}\left[c_3\left(\partial_\vartheta S_0\right)^2-c_2\left(\partial_\vartheta S_0\right)\left(\partial_\zeta S_0\right)\right]+\widetilde{E}^{-1}\left[c_3\left(\partial_\phi S_0\right)^2-c_4\left(\partial_\phi S\right)\left(\partial_\vartheta S\right)\right] \tag{29}$$
$$+c_3m^2=0,$$

$$\widetilde{A}^{-1}\left[c_0\left(\partial_\phi S_0\right)\left(\partial_t S_0\right)-c_4\left(\partial_t S_0\right)^2\right]+\widetilde{B}\left[c_4\left(\partial_r S_0\right)^2-c_1\left(\partial_r S_0\right)\left(\partial_\phi S_0\right)\right]$$
$$+\widetilde{C}^{-1}\left[c_4\left(\partial_\zeta S_0\right)^2-c_2\left(\partial_\zeta S_0\right)\left(\partial_\phi S_0\right)\right]+\widetilde{D}^{-1}\left[c_4\left(\partial_\vartheta S_0\right)^2-c_3\left(\partial_\vartheta S_0\right)\left(\partial_\phi S_0\right)\right] \tag{30}$$
$$+c_4m^2=0.$$

Considering property of spacetime, we carry out the separation of variables as

$$S_0=-\omega t+W\left(r\right)+\Theta\left(\zeta,\vartheta\right)+j\phi, \tag{31}$$

where ω and j are the energy and angular momentum of the emitting particles. Putting Eq. (31) into Eqs. (26) - (30), one obtains a matrix equation $\Lambda\left(c_0, c_1, c_2, c_3, c_4\right)^T$. The components of 5×5 matrix Λ are

$$\begin{aligned}
&\Lambda_{00}=m^2+\widetilde{B}\left(W'\right)^2+\widetilde{C}^{-1}\left(\partial_\zeta\Theta\right)^2+\widetilde{D}^{-1}\left(\partial_\vartheta\Theta\right)^2\\
&+\widetilde{E}^{-1}j^2,\quad \Lambda_{01}=\widetilde{B}W'\omega,\quad \Lambda_{02}=\widetilde{C}^{-1}\omega\left(\partial_\zeta\Theta\right),\\
&\Lambda_{03}=D^{-1}\omega\left(\partial_\vartheta\Theta\right),\quad \Lambda_{04}=E^{-1}\omega j,\\
&\Lambda_{10}=-\widetilde{A}^{-1}\omega W',\\
&\Lambda_{11}=m^2-\widetilde{A}^{-1}\omega^2+\widetilde{C}^{-1}\left(\partial_\zeta\Theta\right)^2\\
&+\widetilde{D}^{-1}\left(\partial_\vartheta\Theta\right)^2+\widetilde{E}^{-1}j^2,\Lambda_{12}=-\widetilde{C}^{-1}W'\left(\partial_\zeta\Theta\right),\\
&\Lambda_{13}=-\widetilde{D}^{-1}\left(\partial_\vartheta\Theta\right)W',\Lambda_{14}=-\widetilde{E}^{-1}jW',\\
&\Lambda_{20}=-\widetilde{A}^{-1}\omega\left(\partial_\zeta\Theta\right),\Lambda_{21}=-\widetilde{B}W'\left(\partial_\zeta\Theta\right),\\
&\Lambda_{22}=m^2-\widetilde{A}^{-1}\omega^2+\widetilde{B}\left(W'\right)^2+\widetilde{D}^{-1}\left(\partial_\vartheta\Theta\right)^2\\
&+\widetilde{E}^{-1}j^2,\Lambda_{23}=-\widetilde{D}^{-1}\left(\partial_\vartheta\Theta\right)\left(\partial_\zeta\Theta\right),\\
&\Lambda_{24}=-\widetilde{E}^{-1}j\left(\partial_\zeta\Theta\right),\Lambda_{30}=-\widetilde{A}^{-1}\omega\left(\partial_\vartheta\Theta\right),\\
&\Lambda_{31}=-\widetilde{B}W'\left(\partial_\vartheta\Theta\right),\Lambda_{32}=-\widetilde{C}^{-1}\left(\partial_\vartheta\Theta\right)\left(\partial_\zeta\Theta\right),\\
&\Lambda_{33}=m^2-\widetilde{A}^{-1}\omega^2+\widetilde{B}\left(W'\right)^2+\widetilde{C}^{-1}\left(\partial_\vartheta\Theta\right)^2\\
&+E^{-1}j^2,\quad \Lambda_{34}=-E^{-1}\left(\partial_\vartheta\Theta\right)j,\\
&\Lambda_{40}=-\widetilde{A}^{-1}\omega j,\quad \Lambda_{41}=-\widetilde{B}W'j,\\
&\Lambda_{42}=-\widetilde{C}^{-1}\left(\partial_\zeta\Theta\right)j,\Lambda_{43}=-\widetilde{D}^{-1}j\left(\partial_\vartheta\Theta\right),\\
&\Lambda_{44}=m^2-\widetilde{A}^{-1}\omega^2+\widetilde{B}\left(W'\right)^2+\widetilde{C}^{-1}\left(\partial_\zeta\Theta\right)^2\\
&+\widetilde{D}^{-1}\left(\partial_\vartheta\Theta\right)^2,
\end{aligned}\tag{32}$$

where $W'=\partial_r S_0$ and $j=\partial_\phi S_0$. For obtaining a nontrivial solution, the determinant of the matrix Λ must equals to zero. Hence,

$$\mathrm{Im}\tilde{W}_\pm=\pm\int\sqrt{\frac{\tilde{C}\tilde{D}\tilde{E}\omega^2+\tilde{X}}{\tilde{A}\tilde{B}\tilde{C}\tilde{D}\tilde{E}}}dr=\pm\pi r_H\omega/2,\tag{33}$$

where $\tilde{X}=-\tilde{A}\tilde{D}\tilde{E}\left(\partial_\zeta\Theta\right)^2-\tilde{A}\tilde{C}\tilde{E}\left(\partial_\vartheta\Theta\right)^2-\tilde{A}\tilde{C}\tilde{D}j^2-\tilde{A}\tilde{C}\tilde{D}\tilde{E}m^2$. The plus (minus) sign corresponds to the outgoing (incoming) solutions of vector particles. As a result, the tunneling rate is

$$\tilde{\Gamma}=\frac{\tilde{\Gamma}_{(emission)}}{\tilde{\Gamma}_{(absorpation)}}=e^{-4\mathrm{Im}W_+}=e^{-2\pi r_H\omega}.\tag{34}$$

With the formula $\Gamma=\exp(E/T)$, where E and T are the energy of emitting particle and the temperature, we can calculate the Hawking temperature of 5-dimensional ST black hole

$$\widetilde{T_H}=\frac{1}{2\pi r_H}=\frac{1}{2\pi\sqrt{M}}.\tag{35}$$

The Hawking temperature of the 5-dimensional ST black hole is only related to the mass. People can get the same result when they investigate the scalar particles tunneling and Dirac particles tunneling from the 5-dimensional ST black hole. When assuming $D = 4$, we find that the Hawking temperature of 4-dimensional ST black hole is $T_{H(4D-ST)} = 1/4\pi r_{H(4D-ST)}$ with $r_{H(4D-ST)} = 2M$, which is different from Eq. (35). This difference is caused by the property of ST black hole spacetime, it indicates that people may obtain different information from higher dimensional black hole.

4 Discussion and conclusion

In this paper, we studied the vector particles tunneling from 4-dimensional KdS black hole and 5-dimensional ST black hole. The tunneling rates and Hawking temperatures were gotten. For the 4-dimensional KdS black hole, we found that the tunneling rates and Hawking temperatures are not only depended on the outer event horizon, mass and angular momentum of the 4-dimensional KdS black hole but also the cosmological constant. Besides, Eq. (21) and Eq. (22) are consistent with that obtained by scalar particles and Dirac particles tunneling from the 4-dimensional KdS black hole. In the static limit, Eq. (22) is reduced to the temperature of SC black hole. For 5-dimensional ST black hole, the $\widetilde{\Gamma}$ and $\widetilde{T_H}$ are related to the mass of the black hole. Moreover, since the $f(r)$ and the event horizon of ST black hole are depended on the dimension of spacetime, they lead the temperature of 5-dimensional ST black hole is different from the temperature of 4-dimensional ST black hole.

Moreover, by applying the WKB approximation, we derived the Hamilton-Jacobi equation from Proca equation. Yang and Lin (2010) and Mu et al. (2015) also derived the Hamilton-Jacobi equation from Klein-Gordon equation, Dirac equation and Rarita-Schwinger equation. Therefore, we think the Hamilton-Jacobi is a fundamental equation in the semiclassical theory, which can help people to investigate the semiclassical Hawking radiation behavior.

The Eq. (22) and Eq. (35) showed the Hawking radiation is the black-body radiation, it indicates that the black holes will emit away all their particles as the black-body radiation, that is, the black hole lose all its information. In order to solve this problem, the results need to be modified. In our further work, we will take into account the conservation of energy and the self-gravitational interaction.

Acknowledgements

This work is supported by the Natural Science Foundation of China (Grant No. 11573022 and No. 11205125), the Sichuan Province Science Foundation for Youths (Grant No. 2014JQ0040), and the Innovative Research Team in China West Normal University (Grant No. 438061).

References

Bekenstein, J. D., 1973, Phys. Rev. D, 7, 2333

Hawking, S. W., 1975, Commun. Math. Phys., 43, 199

Parikh, M. K., Wilczek, F., 2000, Phys. Rev. Lett., 85, 5042

Kerner, R., Mann, R. B., 2008, Class. Quant. Grav, 25, 095014

Kerner, R., Mann, R. B., 2008, Phys. Lett. B, 665, 277

Wang, M. J., Ding, C. K., Chen, S. B., Jing, J. L., 2010, Gen. Relativ. Gravit., 42, 347

Chatterjee, B., Mitra, P., 2009, Phys. Lett. B, 675, 240

Yale, A., 2011, Phys. Lett. B, 697, 398

Yale, A., 2011, arXiv:0805.2220 [hep-th]

Feng, Z. W., Deng, J., Li, G. P., Yang, S. Z., 2012, Int. J. Theor. Phys., 51, 3214

Jiang, Q. Q. Wu, S. Q. Cai, X., 2006, Phys. Rev. D, 73, 064003

Lin, K. Yang, S. Z., 2014, Int. J. Theor. Phys., 53, 1710

Chen, D. Y. Jiang, Q. Q. Wang, P. Yang, H. T., 2013, J. High. Energy. Phys., 1311, 176

Chen, D. Y. Jiang, Q. Q. Zu, X. T., 2008, Phys. Lett. B, 665, 106

Chen, D. Y Li, Z. H., 2014, Adv. High. Energy. Phys., 2014, 620157

Zhang, J. Y Zhao, Z., 2005, Adv. High. Energy. Phys., 20, 1673

Jiang, Q. Q., 2008, Phys. Lett. B, 666, 517

Ding, C. K., Jing, J. L., 2009, Phys. Lett. B, 676, 99

Zhang, J. Y., Zhao, Z., 2005, Phys. Lett. B, 618, 14

Zhang, J. Y. Zhao, Z., 2005, J. High. Energy. Phys., 0505, 055

Jiang, Q. Q., 2012, Commun. Theor. Phys., 58, 237

Li, H. L., Lin, R., Cheng, L. Y., 2012, Europhys. Lett., 98, 30002

Li, H. L., Lin, R., 2011, Gen. Relativ. Gravit., 43, 2115

Kruglov, S. I., 2014, Mod. Phys. Lett. A, 29, 1450203

Kruglov, S. I., 2014, Int. J. Mod. Phys. A, 29, 1450118

Chen, G. R., Zhou, S. W., Huang, Y. C., 2015, Int. J. Mod. Phys. D, 24, 1550005

Carter, B., 1968, Commun. Math. Phys., 10, 280

Feng, Z. W., Zhang, L., Zu, X. T., 2014, Mod. Phys. Lett. A, 29, 1450123

Zheng, Z. M., 2008, Phys. Lett. B, 666, 376

Khani, F., Darvishi, M. T., Baghbani, R., 2013, Astrophys. Space. Sci., 348, 189

Tangherlini, F., 1963, Nuovo Cimento, 27, 636

Abdolrahimi, S., Shoom, A. A., Page, D. N., 2010, Phys. Rev. D, 82, 124039

Beach, S. J. M., Poisson, E., Nickel, B. G., 2014, Phys. Rev. D, 89, 124014

Yang, S. Z., Lin, K., 2010, Sci. Sin. Phys, Mech & Astron, 8240, 507

Jiang, Q. Q., 2007, Class. Quant. Grav., 24, 4391

Wang, S., Wu, S. Q., Xie, F., Dan, L., 2006, Chin. Phys. Lett., 10, 1096

Ali, A. F., Faizal, M., Khalil, M. M., 2015, Nucl.Phys. B, 894, 341

Kruglov, S. I., 2011, Int. J. Mod. Phys. A, 26, 2487

Angheben, M. Nadalini, M. Vanzo, L. Zerbini, Z., 2005, J. High. Energy. Phys., 0505, 014

Shankaranarayanan, S., 2005, Phys. Rev. D, 67, 084026

Mu, B. R., Wang, P., Yang, H. T., 2015, Adv. High. Energy. Phys., 2015, 898916

Ding, C. K., Jing, J. L., 2010, Class. Quant. Grav., 27, 035004